

Beyond Horizon TMUA Mock Paper

Set 4 Paper 1

75 Minutes

20 Questions

One answer per question

1) Find the area bounded by the straight lines $x = -1$ and $x = 1$ and the graphs of $f(x)$ and $g(x)$, where $f(x) = x^3$ and

$$g(x) = \begin{cases} x^5 & \text{if } -1 \leq x \leq 0, \\ x & \text{if } 0 \leq x \leq 1, \end{cases}$$

(A) $\frac{1}{3}$

(B) $\frac{1}{8}$

(C) $\frac{1}{2}$

(D) $\frac{1}{4}$

(E) $\frac{1}{5}$

(F) $\frac{1}{6}$

2) The largest volume of a cube that can be enclosed in a sphere of diameter 2 cm, is, in cm^3 ,

(A) 1

(B) $2\sqrt{2}$

(C) π

(D) $\frac{8}{3\sqrt{3}}$

(E) $\frac{8}{9}$

(F) $\frac{2\sqrt{2}}{3}$

3) If α, β are the roots of $ax^2 + bx + c = 0$, then the equation whose roots are α^2, β^2 is

(A) $a^2x^2 + (b^2 - 2ac)x + c^2 = 0$

(B) $a^2x^2 - (b^2 + 2ac)x + c^2 = 0$

(C) $a^2x^2 - (b^2 - 2ac)x + c^2 = 0$

(D) $a^2x^2 - (a^2 - 2bc)x + c^2 = 0$

(E) $b^2x^2 - (b^2 - 2ac)x + c^2 = 0$

(F) none of the above equations

4) Evaluate the following sum

$$(1 \cdot 1!) + (2 \cdot 2!) + (3 \cdot 3!) + \cdots + (50 \cdot 50!)$$

(A) $51!$

(B) $2 \cdot 5!$

(C) $51! - 1$

(D) $51! + 1$

(E) $50! + 1$

(F) $50! + 1$

5) The number of distinct points common to the curves $x^2 + 4y^2 = 1$ and $4x^2 + y^2 = 4$ is

(A) 0

(B) 1

(C) 2

(D) 3

(E) 4

(F) 5

6) In a parallelogram $PQRS$, P is the point $(-1, -1)$, Q is $(8, 0)$ and R is $(7, 5)$. Then S is the point

- (A) $(-1, 4)$
- (B) $(-2, 4)$
- (C) $(-2, \frac{7}{2})$
- (D) $(-\frac{3}{2}, 4)$
- (E) $(-2, 5)$
- (F) $(-1, 5)$

7) How many real solutions are there to the equation

$$2 \tan^4 x - 5 \tan^2 x + 3 = 0$$

in the interval $0 \leq x < 2\pi$?

- (A) 0
- (B) 1
- (C) 2
- (D) 3
- (E) 4
- (F) 5
- (G) 6
- (H) 7
- (I) 8

8) Given that

$$\frac{dy}{dx} = 4x^2 - \frac{3-2x}{x^3}, \quad x \neq 0$$

and $y = 6$ when $x = 1$, find y in terms of x .

(A) $y = \frac{4}{3}x^3 + \frac{3}{2}x^{-2} - \frac{2}{3}x^{-1} + \frac{31}{6}$

(B) $y = 2x^3 + x^{-2} - x^{-1} + 5$

(C) $y = \frac{4}{3}x^3 + \frac{3}{2}x^{-2} - 2x^{-3} + \frac{23}{6}$

(D) $y = \frac{4}{3}x^3 + \frac{3}{2}x^{-2} - 2x^{-1} + \frac{31}{6}$

(E) $y = \frac{4}{3}x^3 + \frac{3}{2}x^{-2} - \frac{2}{3}x^{-3} + 4$

(F) $y = 3x^3 + x^{-2} - x^{-1} + 3$

9) Evaluate

$$\sum_{n=1}^{50} \log_{10} (2^{1-n})$$

- (A) $-495 \log_{10} 2$
- (B) $495 \log_{10} 2$
- (C) $-1225 \log_{10} 2$
- (D) $1225 \log_{10} 2$
- (E) $1 - 495 \log_{10} 2$
- (F) $1 + 495 \log_{10} 2$
- (G) $1 - 1225 \log_{10} 2$
- (H) $1 + 1225 \log_{10} 2$

10) In the expansion of $(a+bx)^4$ the coefficient of x^3 is 4 times the coefficient of x^2 . Given that a and b are non-zero positive integers, what is the smallest possible value of $a+b$?

- (A) 5
- (B) 6
- (C) 7
- (D) 11
- (E) 15
- (F) 19

11) The functions f and g are given by $f(x) = 4x^2 + 10x + 5$ and $g(x) = x^3 + 5x^2 + 8x - 10$.

What is the complete set of values of x for which one of the functions is increasing and the other decreasing?

- (A) $x \geq -2$
- (B) $x \leq -2$
- (C) $-4 \leq x \leq -3, x \geq -2$
- (D) $x \leq -3, x \geq -2$
- (E) $x \leq -4, -3 \leq x \leq -2$
- (F) $x \leq -4, x \geq -3$
- (G) $-3 \leq x \leq -2$

12) Find the value of

$$\int_1^5 \frac{4-3x}{x\sqrt{x}} dx$$

(A) $\frac{14\sqrt{5}-38}{\sqrt{5}}$

(B) $-\frac{95}{16}$

(C) $\frac{15\sqrt{5}-38}{\sqrt{5}}$

(D) -2

(E) $-\frac{1}{2}$

(F) $\frac{9}{4}$

(G) $\frac{14\sqrt{5}-38}{\sqrt{3}}$

13) Find the number of solutions of the equation

$$x \sin 3x = \cos x$$

in the interval $0 \leq x \leq 3\pi$.

- (A) 1
- (B) 3
- (C) 5
- (D) 7
- (E) 9

14) The number of ways in which three non-negative integers n_1, n_2, n_3 can be chosen such that $n_1 + n_2 + n_3 = 10$ is

(A) 66

(B) 55

(C) 10^3

(D) $\frac{10!}{3!2!1!}$

(E) $\frac{9!}{3!2!1!}$

(F) 10^2

15) Find the complete set of values of the constant c for which the cubic equation

$$x^3 - 3x + c = 0$$

has three distinct real solutions.

(A) $0 < c < 2$

(B) $-2 < c < 0$

(C) $-2 < c < 2$

(D) $c > -2$

(E) $c < 2$

(F) $c < 0$

16) In the interval $0 \leq x \leq 2\pi$, the equation

$$\sin(3 \cos(3x) + 1) = 0$$

has exactly

- (A) 2 solutions
- (B) 4 solutions
- (C) 6 solutions
- (D) 8 solutions
- (E) 10 solutions
- (F) 12 solutions

17) An equilateral triangle has centre O and side length 2. A straight line through O intersects the triangle at two distinct points P and Q . The minimum possible length of PQ is

(A) $\frac{2}{3}$

(B) $\frac{1}{2}$

(C) $\frac{2\sqrt{3}}{3}$

(D) $\frac{4}{3}$

(E) $\sqrt{3}$

(F) $\sqrt{2}$

18) What is the area of the polygon formed by all the points (x, y) in the plane satisfying the inequality

$$|x - 2| + |y - 2| \leq 8?$$

- (A) 32
- (B) 64
- (C) 96
- (D) 128
- (E) 160

19) Let A be a point on the circle with the equation $x^2 + y^2 = 4$, and let B be a point on the circle with the equation $x^2 + y^2 = 9$. If the angle between the vectors representing the points A and B is 30° , what is the distance between the points A and B ?

- (A) $\sqrt{13}$
- (B) $6\sqrt{3}$
- (C) $\sqrt{3}$
- (D) $\sqrt{13 - 6\sqrt{3}}$
- (E) $\sqrt{13 + 6\sqrt{3}}$

20) What is the degree of the following polynomial

$$\frac{d^2}{dx^2} [(3x - 2)^4(2 - x)^5] + \frac{d}{dx} [(3x + 2)^4(4x^2 - 3)^2]$$

- (A) 11
- (B) 10
- (C) 9
- (D) 8
- (E) 7
- (F) less than 7